arXiv:1303.1570v2 [cond-mat.mes-hall] 15 Mar 2013

Localized many particle Majorana fermions with vanishing time reversal symmetry breaking in double quantum dots

Anthony R. Wright*
School of Mathematics and Physics, University of Queensland, Brisbane, 4072 Queensland, Australia

Menno Veldhorst†
ARC Centre of Excellence for Quantum Computation and Communication Technology,
School of Electrical Engineering & Telecommunications,
The University of New South Wales, Sydney 2052, Australia

(Dated: March 18, 2013)

We introduce the concept of spinful many-particle Majorana fermions. These emergent particles are nonlocal in even operator products, but local in odd operator products, thereby preserving their local statistics. We consider a superconductor - double quantum dot system where these modes can arise in the presence of vanishing Zeeman splitting. We find a reverse Mott-insulator transition, where the even parity band crosses the odd parity band. Above this transition, Majorana operators move the system between the odd parity ground state, associated with elastic co-tunneling, and the even parity ground state, associated with crossed Andreev reflection. These Majorana modes are described in terms of one, three and five operator products. Parity conservation results in a 4π periodic supercurrent in the even state and no supercurrent in the odd state. The necessity of only small magnetic fields and the broad parameter space where Majorana fermions appear simplifies the experimental realization greatly, paving the way towards the detection of the Majorana fermion.

The prediction for the existence of Majorana fermions [1] has attracted enormous attention in condensed matter physics in recent years [2–7]. Allured by the possibility of constructing topological qubits for quantum computation [2, 4], a plethora of schemes promising the positive identification of Majorana fermions has emerged [8–12]. Superconductors provide intrinsic electron-hole coupling resulting in chargeless quasiparticles, and most proposals to realize Majorana fermions are based on this principle. Standard s-wave superconductors do not support a zero energy Majorana mode. In the presence of strong Rashba spin-orbit coupling, however, an effective spinless p-wave superconductor can arise. This idea was first recognized by Fu and Kane, who considered the interface of a superconductor and a topological insulator [5]. A topological state harnessing Majorana modes can also arise using semiconducting nanowires in the presence of Zeeman and Rashba fields [6, 7]. Experimentally, supercurrents [13], Fraunhofer patterns and Shapiro steps [14], SQUIDs [15], and zero bias conductance peaks [16] have been observed in topological insulator systems, which together with the zero bias conductance peaks in nanowire systems [17] provide prospects for the observation of the Majorana fermion, but to date, no conclusive evidence has been observed.



FIG. 1: (a) Schematic representation of the device considered. Two superconductors (with phase difference ϕ_-) are connected via a double quantum dot with on-site Coulomb repulsion U and in the presence of a magnetic field $E_{Z,\perp} \sim k_B T$. A nanomagnet introduces a localized $E_{Z,\parallel}$, which rotates the field near one of the two dots, and we define θ to be the relative angle between the local fields. (b) Tuning to the degeneracy of the ground state of the double quantum dot system. The color scale represents the energy gap between the lowest even and odd energy eigenstates, $E_g = (\epsilon_1 - \epsilon_2)/\Delta$. The parameters are $U = 3|\Delta| = -3t$. The degenerate ground states support many-particle Majorana fermions with local statistics, for *all* $\theta \neq 0$. The system is symmetric around $\phi = \pi$, but not around $\theta = \pi/2$, since the presence of regular Andreev reflection distinquishes the quantum dots having parallel and antiparallel spin axes.

Quantum dots can also be used to realize localized Majorana modes [11, 18, 19]. Quantum dot proposals include the use of Rashba coupling [11], but the presence of an anisotropic magnetic field can also mimic this effect [19]. Although these proposals are already experimentally quite promising, the strong Rashba or anisotropic magnetic fields required makes them very challenging and limits material flexibility. For example, anisotropic magnetic fields only result in spinless localized Majorana modes in fields $E_Z \gg \Delta$, with Δ the induced superconducting gap. Nonetheless, here we will show that in the presence of small anisotropic magnetic fields, $|B| \sim k_B T$, a new arena emerges: the concept of *spinful many-particle Majorana fermions*. These Majorana fermions are still localized in their odd operator products, and thereby preserve the local statistics of spinless proposals [21].

We consider an s-wave superconductor - double quantum dot system as depicted in Fig.[1], where crossed Andreev reflection (CAR) dominates elastic co-tunneling (EC), a regime that is readily achieved [22–26]. Double electron occupancy in a quantum dot, as we will show, inhibits, but does not prevent the appearance of Majorana fermions. On-site Coulomb repulsion U can be used to tune the system from the double occupancy regime, which we define as $U = 0$, towards the single electron regime, $U = \infty$. Upon increasing the Coulomb repulsion, we find a clear phase transition where it becomes possible to tune the even and odd parity states to become degenerate, see Fig.[2]. This is reminiscent of a Mott metal-insulator transition, however in reverse, where the onsite Coulomb repulsion drives the system *toward* having a degenerate ground state. Above this transition, we can always tune towards the degeneracy point using a second superconductor with a phase difference ϕ_-. Many-particle Majorana fermions appear right at the crossing and are described by one, three and five operator products. The even operator products are nonlocal, while the odd operator products are local for all angles θ, the angle between the local fields on the dots.

The requirements of only small fields and the use of quantum dots opens large material flexibillity. For example, the system could be based on III-V heterostructures [27], but also on materials such as silicon and diamond with large fractions of non-magnetic nuclei which increase spin coherence times [28]. In order to elucidate the novel many-particle physics that emerges, we will take all results in the limit $E_Z \to 0$, valid as $T \to 0$, meaning we have spinful, spin rotation symmetric, ground states.

The superconductor - double quantum dot system with onsite Coulomb repulsion in the presence of an anisotropic magnetic field, Fig.[1], is described by the Hamiltionian

$$H = H_S + H_D + H_T + H_Z. \tag{1}$$

The superconducting part, H_S, is

$$H_S = \sum_{i=1}^{2} \sum_{k,\sigma} \epsilon_{k,i,\sigma} c^\dagger_{k,i,\sigma} c_{k,i} + \Delta e^{i\phi_i} c^\dagger_{k,i,\sigma} c^\dagger_{-k,i,\bar{\sigma}}. \tag{2}$$

Here, $\bar{\sigma}$ denotes the spin that is not σ. The double quantum dot is described by, H_D,

$$H_D = \sum_{j=1}^{2} \sum_{\sigma} \epsilon_j c^\dagger_{j,\sigma} c_{j,\sigma} + \sum_j U n_{j\uparrow} n_{j,\downarrow}, \tag{3}$$

with U the onsite Coulomb repulsion, and ϵ_j the onsite energy of the j^{th} dot. The third term introduces the tunneling and is described by, H_T,

$$H_T = \sum_{i,j,\sigma} \Gamma_{ij} c^\dagger_{i,\sigma} c_{j,\sigma}. \tag{4}$$

The overlap integral Γ_{ij} is between the i^{th} quasiparticle in the superconducting lead, and the j^{th} dot. Finally,

$$H_Z = -E_Z \sum_j \big(c^\dagger_{j,\uparrow} c_{j,\uparrow} - c^\dagger_{j,\downarrow} c_{j,\downarrow} \big). \tag{5}$$

The Hamiltonian is quadratic in the leads, and so we can integrate out the superconducting leads, following the procedure of Ref.[29], to obtain the effective Hamiltonian of the double dot-superconductor system.

Before presenting the final effective theory, we first consider the anisotropic magnetic field, which defines the spin axes. The angle θ, the angle between the two local magnetic fields, modify EC and CAR as follows:

$$\begin{aligned} t c^\dagger_{1,\sigma} c_{2,\sigma} &\to\ t\cos(\theta/2) c^\dagger_{1,\sigma} c_{2,\sigma} + \bar{\sigma} t \sin(\theta/2) c^\dagger_{1,\sigma} c_{2,\bar{\sigma}} \\ \Delta c_{1,\sigma} c_{2,\bar{\sigma}} &\to\ \sigma \Delta \sin(\theta/2) c_{1,\sigma} c_{2,\sigma} + \Delta \cos(\theta/2) c_{1,\sigma} c_{2,\bar{\sigma}}, \end{aligned} \tag{6}$$

where $\sigma, \bar{\sigma} = \pm$, and t, Δ are the effective EC hopping and CAR Cooper pairing amplitudes [29]. The final effective Hamiltonian of the dot-superconductor system is

$$\begin{aligned} H_{eff} = &\sum_{j,\sigma} \epsilon_j c^\dagger_{j,\sigma} c_{j,\sigma} + U \sum_j n_{j,\uparrow} n_{j,\downarrow} - E_Z \sum_j \big(n_{j,\uparrow} - n_{j,\downarrow} \big) \\ &+ \sum_\sigma \Big(t\cos(\theta/2) c^\dagger_{1,\sigma} c_{2,\sigma} + \bar{\sigma} t \sin(\theta/2) c^\dagger_{1,\sigma} c_{2,\bar{\sigma}} \\ &+ \Delta e^{i\frac{\phi_+}{2}} \cos\big(\frac{\phi_-}{2}\big) \big(\sigma \sin(\theta/2) c^\dagger_{1,\sigma} c^\dagger_{2,\sigma} + \cos(\theta/2) c^\dagger_{1,\sigma} c^\dagger_{2,\bar{\sigma}} \big) \\ &+ \Delta e^{i\frac{\phi_+}{2}} \cos\big(\frac{\phi_-}{2}\big) \sum_j c^\dagger_{j,\sigma} c^\dagger_{j,\bar{\sigma}} + h.c. \Big), \end{aligned} \tag{7}$$

where $\phi_\pm = \phi_1 \pm \phi_2$ is the sum (difference) between the phases of the two superconductors. From hereon we will assume that the onsite energy of the two dots has been tuned to the chemical potential of the superconductors, which we define as our zero of energy ($\epsilon_1 = \epsilon_2 = 0$). A discussion of the effects of the onsite energies deviating from this 'sweet spot' has been presented elsewhere [19].

The Hamiltonian Eq. [7] cannot be decomposed into total spin sectors, as the anisotropic magnetic field mixes these. However, the fermion parity is a conserved quantity. The total Hilbert space has dimension $(2^{n_\sigma})^{n_j}/2 = 8$ for each parity sector (i.e. even and odd), which makes exact diagonalization particularly straight-forward. In the occupation representation, we can define an occupation basis $|1, 2\rangle$, where the numbers correspond to the two dots, and we will use arrows to denote the spin, and then we can construct a basis of the sixteen possible configurations. In the $U \to \infty$ limit, the total Fock space is restricted to nine possible states, 5 for even parity, and 4 for the odd, which can be easily solved to give the eigenstates that determine the ground state, in the even and odd sectors. We solve for finite Zeeman splitting, and obtain in the odd parity sectors that the eigenstates are $\epsilon_{odd} = \pm t \pm E_Z$. We take the limit $E_Z \to 0$ in the lowest energy odd parity state to obtain the limit of vanishingly small magnetic fields of the ground state even and odd parity wavefunctions

$$\begin{aligned} \Psi_{even} = \frac{1}{2} \Big[&\pm\sqrt{2} e^{\frac{i\phi_+}{2}} |0, 0\rangle + \cos(\theta/2) \big(|\uparrow, \downarrow\rangle + |\downarrow, \uparrow\rangle \big) \\ &+ \sin(\theta/2) \big(|\uparrow, \uparrow\rangle - |\downarrow, \downarrow\rangle \big) \Big] \end{aligned} \tag{8}$$



FIG. 2: Phase diagram showing the onset of a degenerate ground state energy gap as a function of on-site Coulomb repulsion, for $|\Delta| = |t|$. Upon increasing the on-site Coulomb repulsion U, a reverse Mott-insulator transition occurs where the lowest energy state changes from even to odd parity. At the crossing point, Majorana operators move the system between the degenerate ground states.

with corresponding eigenvalue $\epsilon_{even} = \mp\sqrt{2}\Delta\cos(\phi_-/2)$, and

$$\Psi_{odd} = \frac{1}{\sqrt{2}}\Big[\sin(\theta/4)\big(|\uparrow,0\rangle + |0,\uparrow\rangle\big) \\ - \cos(\theta/4)\big(|\downarrow,0\rangle - |0,\downarrow\rangle\big)\Big], \tag{9}$$

with eigenvalue $\epsilon_{odd} = -t$ (taking $t > 0$). A degenerate ground state is obtained when $\epsilon_{even} = \epsilon_{odd}$. Crucially, for any $t < \sqrt{2}|\Delta|$, there is always a ϕ_- which can be chosen, depending on the value of θ, such that a degenerate ground state can be obtained.

We emphasize that as we are using the occupation number basis, *the ground state is given by the lowest energy eigenstate*, and not the zero energy eigenstate as in the more familiar Bogolubov de-Gennes theory.

The two degenerate ground states have even and odd parity, respectively, and so are protected from hybridizing when the total system conserves particle number parity. When there are two degenerate ground states, a pair of Majorana operators (γ_1, γ_2) can be constructed which transform the two ground states into each other, such that $\gamma_1\Psi_{odd} = \Psi_{even}$, for example. These Majorana operators are given by

$$\gamma_1^{U\to\infty}(\theta) = \frac{1}{\sqrt{2}}\Bigg[x(\gamma_{1\uparrow} + \gamma_{1\downarrow}) + \Big(\frac{\cos(\theta/2)}{\cos(\theta/4)} - x\Big)n_{2\downarrow}\gamma_{1\uparrow} \\ + \Big(\frac{\sin(\theta/2)}{\sin(\theta/4)} - x\Big)n_{2\uparrow}\gamma_{1\uparrow} + \Big(\frac{\cos(\theta/2)}{\sin(\theta/4)} - x\Big)n_{2\uparrow}\gamma_{1\downarrow} \\ - \Big(\frac{\sin(\theta/2)}{\cos(\theta/4)} + x\Big)n_{2\downarrow}\gamma_{1\downarrow}\Bigg] \tag{10}$$

where $x = (\sin(\theta/4) - \cos(\theta/4))^{-1}$, and

$$\gamma_{1,\uparrow}^{\phi_+} = (e^{i\phi_+/4}c_{1,\uparrow}^\dagger + e^{-i\phi_+/4}c_{1,\uparrow}), \tag{11}$$



FIG. 3: Josephson supercurrent of the even parity sector at $T = 0.01|\Delta|$ for the limiting cases $U = 0$ and ∞ and intermediate $U = 20|\Delta|$, with $|\Delta| = |t|$ and $\theta = \pi/2$. When $U = 0$, regular Andreev reflection dominates and the Josephson current is 2π periodic and no Majorana fermions exist at all. When $U \to \infty$, regular Andreev reflection is completely forbidden, there zero energy states are local odd operator Majorana fermions, and crossed Andreev reflection results in a 4π periodic Josephson effect. For finite U, hybridization between the even parity eigenstates leads to a sharp transition at $\phi = \pi$. For large U, Zener tunneling Γ can restore the 4π periodicity, but Majorana modes are always present. $J_0 = e\Delta/\hbar$ is the maximum supercurrent at $U = 0$. The odd parity supercurrent is always 2π periodic, and small, vanishing completely in the limit $U \to \infty$.

and $n_{i,\sigma} = c_{i,\sigma}^\dagger c_{i,\sigma}$. $\gamma_2^{U\to\infty}(\theta)$ has a similar form with site indices $1, 2$ interchanged. Eq. [11] has the form of a usual Majorana operator [21], except that the phase dependence is the total phase of the two superconductors *divided by four*, or equivalently half the average phase of the two superconductors.

The form of three-operator Majorana fermion in Eq. [10] is nonlocal in general. However, it is interesting to note that the relative phase of the Majorana components, which is responsible in general for their non-Abelian braiding statistics, is still localized, as the nonlocality of the Majoranas as defined above is restricted to number operators only. It is assumed, in the above Majorana expressions, that the system is projected into the singly occupied Fock space, since $U \to \infty$.

When U is finite, a phase transition occurs at a critical value of the on-site repulsion. For Coulomb repulsions below this critical point, the ground state is non-degenerate, with the lowest energy even parity state always having lower energy than the lowest energy odd parity state. In Fig.[2] we have plotted the excitation energy of the first excited state $E_g = (\epsilon_2 - \epsilon_1)/|\Delta|$ as a function of onsite Coulomb repulsion, and relative spin angle θ. The phase transition corresponds to the vanishing of the excitation gap at a critical value of on-site Coulomb repulsion. This critical value varies as a function of angle, but for all angles it is quite small, being of the same order as the effective pairing $|\Delta|$.

For finite on-site Coulomb repulsion, the structure of the ground state, and Majorana states are much more complicated than in the infinite U case, though qualitatively do not differ. The odd parity ground state develops a finite weighting on the terms of the form $|\uparrow\downarrow,\uparrow\rangle$, together with the three equivalent combinations of this, whilst the even parity ground state develops a finite weighting on doubly occupied dots ($|\uparrow\downarrow,0\rangle$ and $|0,\uparrow\downarrow\rangle$), together with the four fermion, double occupied quantum dot pair $|\uparrow\downarrow,\uparrow\downarrow\rangle$. The Majorana fermions, however, retain the same phase locality condition. The Majorana fermions in this case acquire five fermion operator products, together with single and three fermion operator products. Generically, the Majorana has the form

$$\gamma_1 = \gamma_{1,\uparrow}^{\phi_+}\Big(a^\uparrow + \sum_{\sigma j} b^\uparrow_{\sigma j} n_{\sigma j} + \sum_{\sigma\sigma' jj'} c^\uparrow_{\sigma\sigma' jj'} n_{\sigma j} n_{\sigma' j'}\Big) + \uparrow\leftrightarrow\downarrow, \tag{12}$$

where $b^\sigma_{\sigma,1} = c^\sigma_{\sigma,\sigma',1,j'} = c^\sigma_{\sigma',\sigma,j,1} = 0$, and when $\phi = \pi/2$, all operators where $j, j' = 2$ have coefficient zero. The appearance of three and five operator products are a clear generalization of the Majorana fermion concept, which is usually based on single operator products [20, 21].

The regime where Majorana fermions appear can be found by analyzing the Josepshon supercurrent through the double dot system. The Josephson current is calculated from the derivative of the free energy with respect to the superconducting phase difference ϕ_- [30]. In contrast to topological superconductor systems, the odd parity ground states disperse only very weakly with superconducting phase difference, and are strictly 2π periodic. At $U \to \infty$, they are completely flat, as CAR cannot possibly excite the odd parity ground state. *The absence of supercurrent in the odd parity state will be a strong signature of parity conservation.* In Fig.[3] we have plotted the Josephson current for the even parity system as a function of superconducting phase difference for the $U \to \infty$ case, where Majorana fermions are present (i.e. CAR splits the Cooper pairs and a gapless super-current flows between the superconductors) versus the case of regular Andreev reflection where the final term in Eq. [7] dominates, as in the $U = 0$ case. The periodicity of the two cases is clearly visible. We have also plotted an intermediate case, $U = 20|\Delta|$, which is 2π periodic in the adiabatic limit. We note that while in a p-wave superconductor the boundstates are protected by fermion parity, the dispersive bound states in the double quantum dot system have both even parity, and scattering between the branches is possible even when the total parity is conserved. Interestingly, the matrix element coupling the branches is a two Majorana operator product. The anomalous current phase relationship, however, can still be observed in non-equilibrium measurements [31] or using dc SQUIDs [32, 33].

At finite U, the bonding and anti-bonding even parity states hybridize as ϕ_- drives them toward a degeneracy point, naturally resulting in a signal where the bonding state tunnels into the anti-bonding, and one where it does not, as shown in Fig.[4]. A superposition of a 2π periodic signal and a 4π periodic signal is expected in this case, since by means of Zener tunneling Γ the quasiparticle can overcome the hybridization gap [34, 35]. This transition from 4π to 2π periodicity is not a transition where Majorana modes disappear, but a transition where the five operator products vanish. Also, finite U results in a weakly dispersing odd parity ground state, clearly visible in Fig.[4], and introduces a supercurrent in the odd parity state.



FIG. 4: Energy eigenstates of the double dot system for U=20$|\Delta|$, $|\Delta| = |t|$ and $\theta = \pi/2$. The $+,-$ signs indicate the even parity ground state bonding and antibonding state, determined by the relative sign between the empty and two-fermion occupations, for example $\Psi_\pm = a|0000\rangle \pm b|1010\rangle + \ldots$. At infinite U, the two states are orthogonal, but at finite U the two develop an anticrossing. The arrow indicates the evolution of the ground state that gives a 4π periodic Josephson effect, which is still possible for small hybridization gaps by means of Zener tunneling Γ. As the bonding and anti-bonding states become increasingly hybridized, the 2π periodic Josephson effect becomes dominant. We emphasize that as we are using the occupation number basis, the ground state is given by the lowest energy eigenstate, and not the zero energy eigenstate as in the more familiar Bogolubov de-Gennes theory.

In conclusion, we have presented an experimentally feasible scenario whereby many particle Majorana fermions in double quantum dots may be realized. We have shown that a large Zeeman splitting is not necessary to realize nonlocal Majorana fermions in double quantum dots, due to the onsite Coulomb repulsion, and in fact a small Zeeman field is sufficient. Furthermore, we have shown that by connecting two out of phase superconductors to both quantum dots, the parameter space can be tuned so as to obtain a degenerate ground state protected by parity, with Majorana operators connecting the ground state. The Majorana operators are local in the odd operators, which determine their statistics, for any angle θ. The effect of a finite, rather than infinite, on-site interaction does not affect the locality conditions of the Majoranas, nor their parity. We expect that the proposed device will expand Majorana physics even further, not in the least due to its spinful ground state, and the fact that the corresponding Majorana fermions are constructed from one, three, and five fermion creation/annihilation operator products, together with their Hermitian conjugates.

We would like to thank Ross H. McKenzie, Jacopo Sabatini and Andrew Dzurak for enlightening discussions. ARW is financially supported by a University of Queensland Postdoctoral Reasearch Fellowship. MV is financially supported by the Australian Research Council Centre of Excellence for Quantum Computation and Communication Technology (project number CE11E0096), the US Army Research Office (W911NF-13-1-0024) and the Netherlands Organization for Scientific Research (NWO) by a Rubicon grant.

---

[*] Electronic address: a.wright7@uq.edu.au
[†] Electronic address: m.veldhorst@unsw.edu.au

[1] E. Majorana, *Nuovo Cimento* **14**, 171-184 (1937).
[2] G. Moore, and N. Read, Nucl. Phys. **B360**, 362 (1991).
[3] A. Kitaev, Ann. Phys. (N.Y.) **303**, 2 (2003).
[4] C. Nayak , S. H. Simon, A. Stern, M. Freedman, and S. Das Sarma, Rev. Mod. Phys. **80**, 1083 (2008).
[5] L. Fu and C.L. Kane, Phys. Rev. Lett. **100**, 096407 (2008).
[6] J.D. Sau, R.M. Lutchyn, S. Tewari, S.D. Sarma, Phys. Rev. Lett. **104**, 040502 (2010).
[7] J. Alicea, Phys. Rev. B **81**, 125318 (2010).
[8] S. Das Sarma, S. M. Freedman, and C. Nayak, Phys. Rev. Lett. **94**, 166802 (2005).
[9] A. R. Akhmerov, J. Nilsson, and C. W. J. Beenakker, Phys. Rev. Lett. **102**, 216404 (2009).
[10] Y. Oreg, G. Refael, and F. von Oppen, Phys. Rev. Lett. **105**, 177002 (2010).
[11] J. D. Sau and D. Das Sarma, Nature Comm. **3**, 964 (2012).
[12] A. R. Wright and B. Rosenow, Phys. Rev. B **86**, 115329 (2012).
[13] B. Sacépé, J.B. Oostinga, J.L. Li, A. Ubaldini, N.J.G. Couto, E. Giannini, and A.F. Morpurgo, Nature Comm. **2**, 575 (2011).
[14] M. Veldhorst, M. Snelder, M. Hoek, T. Gang, X.L. Wang, V.K. Guduru, U. Zeitler, W.G. v.d.Wiel, A.A. Golubov, H. Hilgenkamp, and A. Brinkman, Nature Materials **11**, 417 (2011).
[15] M. Veldhorst, C.G. Molenaar, X.L. Wang, H. Hilgenkamp, and A. Brinkman, Appl. Phys. Lett. **100**, 072602 (2012).
[16] S. Sasaki, M. Kriener, K. Segawa, K. Yada, Y. Tanaka, M. Sato, and Y. Ando, Phys. Rev. Lett. **107**, 217001 (2011).
[17] V. Mourik, K. Zuo, S.M. Frolov, S.R. Plissard, E.P.A.M. Bakkers, and L.P. Kouwenhoven, Science **336**, 1003 (2012).
[18] I.C. Fulga, A. Haim, A.R. Akhmerov, and Y. Oreg, arXiv:1212.1355 (2012).
[19] M. Leijnse, and K. Flensberg, Phys. Rev. B **86**, 134528 (2012).
[20] C. Nayak and F. Wilczek, Nucl. Phys. **B479**, 529 (1996).
[21] D. A. Ivanov, Phys. Rev. Lett. **86**, 268 (2001).
[22] J.M. Byers and M.E. Flatté, Phys. Rev. Lett. **74**, 306 (1995).
[23] G. Deutscher and D. Feinberg, Appl. Phys. Lett. **76**, 487 (2000).
[24] L. Hofstetter *et al.*, Nature **461**, 960 (2009).
[25] L. G. Herrmann *et al.*, Phys. Rev. Lett. **104**, 026801 (2010).
[26] M. Veldhorst and A. Brinkman, Phys. Rev. Lett. **105**, 107002 (2010).
[27] R. Hanson, L.P. Kouwenhoven, J.R. Petta, S. Tarucha, and L.M.K. Vandersypen, Rev. Mod. Phys. **79**, 1217 (2007).
[28] F. Zwanenburg, A.S. Dzurak, A. Morello, M.Y. Simmons, L.C.L. Hollenberg, G. KLimeck, S. Rogge, S.N. Coppersmith, and M.A. Eriksson arXiv arXiv:1206.5202v1.
[29] A. V. Rozhkov and D. P. Arovas, Phys. Rev. Lett. **82**, 2788 (1999).
[30] S. Droste, S. Andergassen, and J. Splettstoesser, J. Phys.: Condens. Matter **24**, 415301 (2012).
[31] D.M. Badiane, M. Houzet, and J.S. Meyer, Phys. Rev. Lett. **107**, 177002 (2011).
[32] M. Veldhorst, C.G. Molenaar, C.J.M. Verwijs, H. Hilgenkamp, and A. Brinkman, Phys. Rev. B **86**, 024509 (2012).
[33] M. Veldhorst, M. Snelder, M. Hoek, C.G. Molenaar, D.P. Leusink, A. A. Golubov, H. Hilgenkamp, and A. Brinkman, Phys. Stat. Sol. RRL **7**, 26 (2013).
[34] A. Jacobs, R. Kümmel, Phys. Rev. B **71**, 184504 (2005).
[35] H. Kroemer, Superlattices Microstruct. **25**, 877 (1999).